

„Projekt Jungbrunnen“ QUELLEN

- Yoshino, J., Mills, K. F., Yoon, M. J. & Imai, S. Nicotinamide mononucleotide, a key NAD(+) intermediate, treats the pathophysiology of diet- and age-induced [diabetes](#) in mice. *Cell Metab.* 14, 528–536 (2011).
- Gomes, A. P. et al. Declining NAD(+) induces a pseudohypoxic state disrupting nuclear-mitochondrial communication during aging. *Cell* 155, 1624–1638 (2013).
- Braidy, N. et al. Mapping NAD(+) metabolism in the brain of ageing Wistar rats: potential targets for influencing brain senescence. *Biogerontology* 15, 177–198 (2014).
- Ying, 2008 W. Ying NAD⁺/NADH and NADP⁺/NADPH in cellular functions and cell death: regulation and biological consequences *Antioxid. Redox Signal.*, 10 (2008), pp. 179-206
- Blander and Guarente, 2004 G. Blander, L. Guarente The Sir2 family of protein deacetylases *Annu. Rev. Biochem.*, 73 (2004), pp. 417-435
- Hall et al., 2013 J.A. Hall, J.E. Dominy, Y. Lee, P. Puigserver, The sirtuin family's role in aging and age-associated [pathologies](#) *J. Clin. Invest.*, 123 (2013), pp. 973-979
- Haigis and Sinclair, 2010 M.C. Haigis, D.A. Sinclair [Mammalian sirtuins: biological insights and disease relevance](#) *Annu. Rev. Pathol.*, 5 (2010), pp. 253-295
- Gross and Henderson, 1983, C.J. Gross, L.M. Henderson, Digestion and absorption of NAD by the small intestine of the rat *J. Nutr.*, 113 (1983), pp. 412-420
- Trammell, S. A., Yu, L., Redpath, P., Migaud, M. E. & Brenner, C. Nicotinamide riboside is a major NAD⁺ precursor vitamin in cow milk. *J. Nutr.* 146, 957–963 (2016).
- Gazzaniga, F., Stebbins, R., Chang, S. Z., McPeck, M. A. & Brenner, C. Microbial NAD metabolism: lessons from comparative genomics. *Microbiol. Mol. Biol. Rev.* 73, 529–541 (2009).
- Birjmohun et al., 2005, Efficacy and safety of high-density [lipoprotein](#) cholesterol-increasing compounds: a meta-analysis of randomized controlled trials, *J. Am. Coll. Cardiol.*, 45 (2005), pp. 185-197
- Anderson et al., 2003, Nicotinamide and PNC1 govern lifespan extension by calorie restriction in *Saccharomyces cerevisiae*, *Nature*, 423 (2003), pp. 181-185
- Trammell SAI et al. Nicotinamide riboside is uniquely and orally bioavailable in mice and humans, *Nature Communications* volume7, Article number: 12948 (2016).
- Costford et al., 2010. Skeletal muscle NAMPT is induced by exercise in humans, *Am. J. Physiol. Endocrinol. Metab.*, 298 (2010), pp. E117-E126
- Chen et al., 2008, Tissue-specific regulation of SIRT1 by calorie restriction, *Genes Dev.*, 22 (2008), pp. 1753-1757
- Yoshino et al., 2011 J. Yoshino, K.F. Mills, M.J. Yoon, S. Imai, Nicotinamide mononucleotide, a key NAD(+) intermediate, treats the pathophysiology of diet- and age-induced diabetes in mice, *Cell Metab.*, 14 (2011), pp. 528-536
- Kraus et al., 2014 Nicotinamide N-methyltransferase knockdown protects against diet-induced obesity, *Nature*, 508 (2014), pp. 258-26
- Schmeisser K, Mansfeld J, Kuhlow D, Weimer S, Priebe S, Heiland I, Birringer M, Groth M, Segref A, Kanfi Y, Price NL, Schmeisser S, Schuster S, Pfeiffer AF, Guthke R, Platzer M, Hoppe T, Cohen HY, Zarse K, Sinclair DA, Ristow M. Role of sirtuins in lifespan regulation is linked to methylation of nicotinamide. *Nat Chem Biol.* 2013 Nov;9(11):693-700. doi: 10.1038/nchembio.1352. Epub 2013 Sep 29. PubMed PMID: 24077178
- Hashimoto T, Horikawa M, Nomura T, Sakamoto K. Nicotinamide adenine dinucleotide extends the lifespan of *Caenorhabditis elegans* mediated by sir-2.1 and daf-16. *Biogerontology.* 2010 Feb;11(1):31-43. doi: 10.1007/s10522-009-9225-3. Epub 2009 Apr 16. PubMed PMID: 19370397.